

Rescon DM Liquid

Drug Facts

Active Ingredients Purpose

Chlorpheniramine Maleate 2 mg...Antihistamine

Dextromethorphan Hydrobromide 10 mg .. Antitusive

d-Pseudoephedrine Hydrochloride 30 mg..Nasal decongestant in each 5 ml of clear, sweet-tasting liquid

Uses: temporarily relieves these symptoms due to cold and flu or allergic reaction:

- cough
- nasal congestion
- runny nose
- sneezing
- itching of nose or throat

Warnings -

Ask a doctor before use if you have:

- persistent or chronic cough
- fever
- rash
- persistent headache
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- emphysema
- glaucoma
- chronic bronchitis
- enlarged prostate gland

Do not use this product if you have taken a monoamine oxidase inhibitor (MAOI) within the last two weeks. If you are uncertain whether your prescription drug contains a MAOI, consult your doctor.

When using this product:

- you may get drowsy
- sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if nervousness, dizziness, or sleeplessness occur. If pregnant or breastfeeding, ask a health professional before use. Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

Adults and children 12 years and older: 2 teaspoonfuls every 4 to 6 hours; Children 6 years to under 12 years: 1 teaspoonful every 4 to 6 hours; Children under 6: Ask your doctor. Do not exceed 4 doses in a 24-hour period.

Other information: store at 15C- 30C (59F-86F)

Inactive ingredients: cherry flavoring, methylparaben, polyethylene glycol, propylparaben, saccharin sodium, sorbitol, and water. This product is dye-free.

Manufactured for
Capellon Pharmaceuticals, Ltd.
Fort Worth, TX 76118
NDC 64543-105-04